

# Predictive value of group I oral lesions in detecting HIV infection amongst patients attending PHC facilities in Gauteng<sup>i</sup>

Bhayat A

A research report submitted to the Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, in partial fulfilment of the requirements for the degree of Master of Dentistry in the branch of Community Dentistry.

JANUARY 2007

## Declaration

I, Ahmed Bhayat, hereby declare that this research report is my own work. It is being submitted for the degree of Master of Dentistry in Community Dentistry in the University of the Witwatersrand, Johannesburg. It has not been submitted or presented for any degree or examination at this or any other University.

.....

.....day of March 2007.

## Dedication

To my family, friends and colleagues for their help and support.

## Publications and presentations

1. Presented at the Prakash Vallabh Primary Health Care (PHC) Conference in Pretoria 2006. Won first prize in the “Academic” category.
2. Presented at the International Association of Dental Research (IADR) conference in Midrand 2006.

## Abstract

The utilization of oral lesions as a screening tool for HIV is not well documented. Attendees at two primary health care facilities (Khutsong and Heidelberg) were assessed to determine the predictive value of group I oral lesions for HIV infection. The objectives were to investigate the: 1) HIV prevalence amongst attendees at PHC facilities, 2) Prevalence of HIV-related oral lesions and 3) Correlation between the oral lesions and the HIV status using the Likelihood Ratio test.

**Methods:** All patients over 12 months of age presenting at the two facilities for a curative care consultation over a one-week period (in April 2005) were included. Consent was obtained by trained counselors who also conducted a brief interview and offered pre-test counseling to patients wishing to know their HIV status. Two calibrated dentists conducted a head, neck and oral examination and administered a rapid saliva HIV test (OraQuick HIV-1/2-Rapid HIV-1/2 Antibody Test).

**Results:** A total of 654 attendees were surveyed in the 2 facilities. There was a 100% response. The mean age of the participants was 34 years (range: 1-94), and the majority (73%) were female. HIV prevalence rates were 34% at Khutsong and 36% at Heidelberg. The HIV prevalence peaked at 46% in the 16-45 age groups. Of the 228 who tested positive for HIV, 121 (53%) patients were diagnosed with 1 or more Group I oral lesion. Oral candidiasis (46%) and oral hairy leukoplakia (19%) were the two most common oral lesions diagnosed in the HIV positive cohort. The positive predictive values and specificity values for multiple lesions ranged between 96% and 100%. Most of the likelihood ratios for multiple lesions were greater than 10 which implied that the patients who presented with these lesions were extremely likely to test positive for HIV. The sensitivity values (1% to 37%) and negative predictive values (66% to 70%) remained relatively low.

**Conclusion:** The HIV prevalence of patients attending PHC facilities was high (34%). Oral lesions are useful markers of HIV-infection and should alert clinicians to the presence of HIV infection. Multiple group I lesions were more predictive of HIV infection compared to single lesions.

## Acknowledgements

I would like to acknowledge the following people who made this report possible:

- Prof Rudolph and Dr. Yengopal for their support and supervision throughout the duration of this project
- Prof Schneider who had the confidence in us to carry out the research
- Ms S Oyedele who helped obtain the data and made sure everything always ran as smoothly as possible

# TABLE OF CONTENTS

	<b>Page</b>
DECLARATION	ii
DEDICATION	iii
PUBLICATIONS AND PRESENTATIONS	iii
ABSTRACT	iv
ACKNOWLEDGEMENTS	v
TABLE OF CONTENTS	vi
LIST OF FIGURES	viii
LIST OF TABLES	ix

## **Chapter 1**

1.0 Introduction	1
1.1 Literature review	2
1.2 Aim	17
1.3 Objectives	17

## **Chapter 2**

2.0 Study design	18
2.1 Study population	19
2.2 Study sample	20
2.3 Informed consent	20
2.4 Measurement	21
2.5 Tests used to determine the predictive value of oral lesions	24
2.6 Ethical clearance	26
2.7 Data collection	26
2.8 Data analysis and statistical tests used	26
2.9 Limitations	27

<b>Chapter 3</b>	
3.0 Response rate	28
3.1 Gender	28
3.2 Age	29
3.3 HIV status	30
3.4 Types of oral lesions diagnosed	31
3.5 The prevalence of oral lesions in relation to the patients' HIV status	33
3.6 Strength of the association between oral lesions and HIV status	33
 <b>Chapter 4</b>	
Discussion	37
 <b>Chapter 5</b>	
Conclusions and recommendations	49
 <b>REFERENCES</b>	51
 <b>APPENDICES</b>	57

## LIST OF FIGURES

<b>Figure</b>		<b>Page</b>
Figure 1	Map of Gauteng Province	19
Figure 2	Interpretation of the saliva test	24
Figure 3	Gender distribution according to the PHC facility	29
Figure 4	The total number of patients per age group per facility	29
Figure 5	The percentage of HIV positive patients classified according to age groups and gender	31



## LIST OF TABLES

Table		Page
Table 1	Studies reporting on the prevalence of group I oral lesions in HIV positive patients	4
Table 2	Results of epidemiological tests performed by Patton (2000)	10
Table 3	Formulae to calculate the Likelihood Ratio	11
Table 4	Formulae to calculate the epidemiological tests	25
Table 5	Daily attendances at the two PHC facilities	28
Table 6	Distribution of patients according to gender and HIV status	30
Table 7	The prevalence of Group I oral lesions in the study population	32
Table 8	The prevalence of oral lesions in relation to the patients' HIV status	33
Table 9	The association between the HIV status and the presence of oral lesions using logistic regression	34
Table 10	Statistical analysis performed on specific oral lesions	36

---